

BCU Mathematics Contest 2000 Solutions

Edited by Pascal Hitzler and Finbarr Holland

March 18, 2000

Department of Mathematics, National University of Ireland — Cork, Cork, Ireland.
E-mail: phitzler@ucc.ie, f.holland@ucc.ie

Contest Web Page:
<http://maths.ucc.ie/~pascal/verein/internat/index.html>

Problems by Pascal Hitzler, Finbarr Holland and Gudrun Kalmbach H. E.

Local Organization:

Bratislava	(Slovak Republic)	Michal Zajac
Cork	(Ireland)	Pascal Hitzler, Finbarr Holland
Ulm	(Germany)	Gudrun Kalmbach H.E.
Wegberg	(Germany)	Franz Josef Klengen

BCU Mathematics Contest, March 18, 2000

INDIVIDUAL CONTEST PAPER Solutions

1 Note that x can not be 0. Multiplying by $|x|$ on both sides of the inequality yields

$$|3x - 2| < 2|x| - 1.$$

We have to distinguish three cases:

a) $x \geq \frac{2}{3}$: Then the inequality yields

$$3x - 2 < 2x - 1$$

from which we obtain that $x < 1$. So all x with $\frac{2}{3} \leq x < 1$ satisfy the inequality.

b) $0 \leq x \leq \frac{2}{3} = \frac{10}{15}$: Then the inequality yields

$$-3x + 2 < 2x - 1$$

and hence $x > \frac{3}{5} = \frac{9}{15}$. So all x with $\frac{3}{5} < x \leq \frac{2}{3}$ satisfy the inequality.

c) $x < 0$: Then the inequality yields

$$-3x + 2 < -2x - 1$$

from which we obtain $3 < x$ which is impossible.

Summing up the three cases, we obtain that exactly all x with

$$\frac{3}{5} < x < 1$$

satisfy the inequality.

2 Denote f by $f^{(0)}$. We obtain for all non-negative integers k :

$$f^{(n)} = 2^n \sin 2x \quad \text{if } n = 4k \quad (1)$$

$$f^{(n)} = 2^n \cos 2x \quad \text{if } n = 4k + 1 \quad (2)$$

$$f^{(n)} = -2^n \sin 2x \quad \text{if } n = 4k + 2 \quad (3)$$

$$f^{(n)} = -2^n \cos 2x \quad \text{if } n = 4k + 3 \quad (4)$$

Intersections with the x -axis:

- (1) $x = \frac{l\pi}{2}$ for all integers l
- (2) $x = \frac{l\pi}{2} + \frac{\pi}{4}$ for all integers l
- (3) $x = \frac{l\pi}{2}$ for all integers l
- (4) $x = \frac{l\pi}{2} + \frac{\pi}{4}$ for all integers l


Maxima:

- (1) $x = l\pi + \frac{\pi}{4}$ for all integers l
- (2) $x = l\pi$ for all integers l
- (3) $x = l\pi - \frac{\pi}{4}$ for all integers l
- (4) $x = l\pi + \frac{\pi}{2}$ for all integers l

Minima:

- (1) $x = l\pi - \frac{\pi}{4}$ for all integers l
- (2) $x = l\pi + \frac{\pi}{2}$ for all integers l
- (3) $x = l\pi + \frac{\pi}{4}$ for all integers l
- (4) $x = l\pi$ for all integers l

Points of Inflection: Exactly the intersections with the x -axis above.



Let H be the mid-point of AB . Then HM_1M_2 are collinear. Since S is the centroid of $\triangle ABE$, HSE are collinear. So HM_2E form a triangle with M_1 on HM_2 and S on HE . Hence M_1E intersects SM_2 .

4 (a) By taking $z = x$ in equation (2) we obtain

$$f(x, x) \leq 2f(x, y)$$

for all x, y , and since $f(x, x) = 0$ by (1), we obtain $f(x, y) \geq 0$.

(b) By taking $y = x$ in equation (2) we obtain

$$f(x, z) \leq f(x, x) + f(z, x) = f(z, x).$$

Since equation (2) holds for all x, y, z , we can also write it as

$$f(z, x) \leq f(z, y) + f(x, y),$$

and by taking $z = y$ we obtain

$$f(z, x) \leq f(z, z) + f(x, z) = f(x, z).$$

Hence $f(x, z) \leq f(z, x)$ and $f(z, x) \leq f(x, z)$ and so $f(z, x) = f(x, z)$.

An example of such a function would be the zero function $f(x, y) = 0$ for all x, y . Indeed, possible examples are exactly all pseudo-metrics on the real numbers.

BCU Mathematics Contest, March 18, 2000


TEAM CONTEST PAPER Solutions

- 1 The area of an equilateral triangle with side-length b is $\frac{\sqrt{3}}{4}b^2$.

The radius of one circle is $r = \frac{\sqrt{3}}{3}a$.

The area of one circle is $A_c = \frac{1}{3}\pi a^2$.

The area of the shamrock is $A = 3A_c - 3B$, where B is the area of the intersection between two circles.




We can calculate

$$\begin{aligned} B &= 2 \left(\frac{1}{6} A_c - \frac{\sqrt{3}}{4} r^2 \right) \\ &= 2 \left(\frac{1}{6} \cdot \frac{1}{3} \pi a^2 - \frac{\sqrt{3}}{4} \left(\frac{\sqrt{3}}{3} a \right)^2 \right) \\ &= \left(\frac{\pi}{9} - \frac{\sqrt{3}}{6} \right) a^2 \\ &= \frac{2\pi - 3 \cdot \sqrt{3}}{18} a^2. \end{aligned}$$

We finally obtain

$$\begin{aligned}
 A &= 3A_c - 3B = \pi a^2 - \left(\frac{\pi}{3} - \frac{\sqrt{3}}{2}\right) a^2 \\
 &= \left(\frac{2\pi}{3} + \frac{\sqrt{3}}{2}\right) a^2
 \end{aligned}$$

2



The easiest way to solve this is to write the curves as functions in y . The first two equalities then yield the function

$$f : y \mapsto y^2 - 4\pi^2$$

and the last equality the function

$$g : y \mapsto \sin y.$$

In order to calculate the intersection points, we have to solve the equality

$$\sin y = y^2 - 4\pi^2. \tag{*}$$

By guessing, we obtain the two solutions $y = \pm 2\pi$.

We next show that $\pm 2\pi$ are the only solutions to the equality (*). Indeed, we know that the slope of the \sin function is at most 1. The slope of the function f at $y = \pm 2\pi$ is

$\pm 4\pi$. Also, $f'(y) < 1 \leq g'(y)$ for $y \leq -1$ and $f'(y) > 1 \geq g'(y)$ for $y \geq +1$. However, for $|y| \leq 1$ we have that $f(y) \leq 1 - 4\pi < -1$, so $\pm 2\pi$ are the only intersection points.

We finally have to solve the integral

$$\int_{-2\pi}^{2\pi} \sin y - y^2 + 4\pi^2 dy$$

which yields

$$\begin{aligned} \left[\cos y - \frac{1}{3}y^3 + 4\pi^2 y \right]_{-2\pi}^{2\pi} &= 1 - \frac{1}{3}8\pi^3 + 8\pi^3 - \left(1 + \frac{1}{3}8\pi^3 - 8\pi^3 \right) \\ &= \frac{4}{3}8\pi^3 = \frac{32}{3}\pi^3. \end{aligned}$$

- 3** (a) By trying, we find solutions for $n = 3, 5, 6, 8, 9, 10$. Now if there is a solution for some n , there obviously is a solution for all $n + 3k$, where k is a positive integer, since we can just add k groups of 3 members each. Since 8, 9, 10 are three consecutive numbers, we obtain as an answer that it is possible for $n = 3, 5, 6$ and for all $n \geq 8$.

(b) We first determine the number $f(n)$ of groups as a function of n . This yields

$$\begin{aligned} f(n) &= \begin{cases} \frac{n}{3} & \text{if } n \text{ is divisible by } 3 \\ \frac{n-10}{3} + 2 & \text{if } n-1 \text{ is divisible by } 3 \text{ and } n \geq 10 \\ \frac{n-5}{3} + 1 & \text{if } n-2 \text{ is divisible by } 3 \end{cases} \\ &= \begin{cases} \frac{n}{3} & \text{if } n \text{ is divisible by } 3 \\ \frac{n-4}{3} & \text{if } n-1 \text{ is divisible by } 3 \text{ and } n \geq 10 \\ \frac{n-2}{3} & \text{if } n-2 \text{ is divisible by } 3 \end{cases} \end{aligned}$$

The function g which is to be determined is then obtained as

$$\begin{aligned} g(n) = \frac{n}{f(n)} &= \begin{cases} 3 & \text{if } n \text{ is divisible by } 3 \\ \frac{\frac{n}{\frac{n-10}{3}+2}}{\frac{n-10}{3}+2} & \text{if } n-1 \text{ is divisible by } 3 \text{ and } n \geq 10 \\ \frac{\frac{n}{\frac{n-5}{3}+1}}{\frac{n-5}{3}+1} & \text{if } n-2 \text{ is divisible by } 3 \end{cases} \\ &= \begin{cases} 3 & \text{if } n \text{ is divisible by } 3 \\ \frac{3n}{n-4} & \text{if } n-1 \text{ is divisible by } 3 \text{ and } n \geq 10 \\ \frac{3n}{n-2} & \text{if } n-2 \text{ is divisible by } 3 \end{cases} \end{aligned}$$

BCU Mathematics Contest, March 18, 2000

SPEED CONTEST PAPER FOR TEAMS Solutions

1 We obtain

$$\begin{aligned}p_0(x) &= x^2 + 1 \\p_1(x) &= x^2 \\p_2(x) &= -x^2 - x^2 - 1 = -2x^2 - 1 \\p_3(x) &= x^2 + x^2 + 1 - x^2 = x^2 + 1 \\p_4(x) &= -x^2 - 1 + 2x^2 + 1 = x^2\end{aligned}$$

and the process repeats. We obtain only three different polynomials.

2 Since the last digit of 1503985847 is a seven, a must be either 1, 3, 7, or 9. The number 1503985847 is not divisible by 3, but for $a = 3$ or 9, $934152a$ is divisible by 3. So a can not be 3 or 9. Trying $a = 1$ we obtain $1503985847 = 9341521 \cdot 161 + 966 = (9341521 + 6) \cdot 161$. Hence a must be 7.

3 The function is

$$f(n) = \begin{cases} n - 10 & \text{if } n > 100 \\ f(f(n + 11)) & \text{if } n \leq 100 \end{cases}$$

(called McCarthy's 91 function). We obtain

$$\begin{aligned}f(98) &= f(f(109)) \\ &= f(99) \\ &= f(f(110)) \\ &= f(100) \\ &= f(f(111)) \\ &= f(101) \\ &= 91.\end{aligned}$$


(Indeed the function is constant 91.)

4 Player B chooses $b = 0$ and $b' = 1$. We obtain the two equations

$$\begin{aligned} ax + y &= c \\ a'y + z &= c' \end{aligned}$$

with the solution $x = 0$, $y = c$ and $z = c' - a'c$.

5 Let the length of DE be x and y be the distance between DE and AB .



Then we obtain

$$\frac{4}{x} = \frac{5}{5-y}.$$

and this yields

$$y = 5 - \frac{5}{4}x.$$

We want to maximise the function f where

$$f(x) = 5x - \frac{5}{4}x^2.$$

We obtain

$$f'(x) = -\frac{5}{2}x + 5$$

which is zero for $x = 2$ only. Since $f''(2) = \frac{-5}{2}$, we obtain a maximum for $x = 2$, so DE is 2cm long.

6

$$x = y = z = \frac{\pi}{2}.$$

Because, in the first place, $1 \geq \sin x, \sin y, \sin z > 0$ and so the product $\sin x \sin y \sin z$ is strictly less than 1 if any factor is less than 1. Hence each factor equals 1.

7 We note that $p(x) = (x - \alpha)(x - \beta)(x - \gamma)$, hence

$$\begin{aligned} p(1) &= (1 - \alpha)(1 - \beta)(1 - \gamma) \\ -p(-1) &= (1 + \alpha)(1 + \beta)(1 + \gamma) \end{aligned}$$

and consequently

$$\frac{1 - \alpha}{1 + \alpha} \cdot \frac{1 - \beta}{1 + \beta} \cdot \frac{1 - \gamma}{1 + \gamma} = \frac{p(1)}{-p(-1)} = -1.$$